



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

where P , Q , and R are numbers which were computed by infinite series (G. Chrystal, *Algebra*, Part II, chapt. 28), it is found, on comparing the results with the known values of these logarithms to 15 decimals, that there are the following errors: $-2,533$, $-4,052$, and $6,080$, respectively. Find the errors in P , Q and R .

GEOMETRY.

When this issue was made up no solutions had been received for numbers 410, 417, 421, and 427. Please give attention to these.

434. Proposed by CLIFFORD N. MILLS, Bloomington, Ind.

ABC is any triangle with sides a , b , c . Prove by purely geometrical methods that the area $A = \sqrt{s(s-a)(s-b)(s-c)}$, where $s = 1/2(a+b+c)$. (From Olney's *Geometry*, page 255, with slightly simplified notation.)

435. Proposed by R. M. MATHEWS, Riverside, California.

From a fixed point A perpendiculars are dropped to the tangents drawn to a circle whose center is O . Prove that the locus of the feet of the perpendiculars is a limaçon.

436. Proposed by A. J. KEMPNER, University of Illinois.

Given in a plane two similar curves arbitrarily situated, except that they shall possess the same sense of direction (which, of course, does not mean that they are similarly located). Let corresponding points on both curves be joined by straight lines, and let all of these straight lines be divided in the same ratio $\lambda : 1$, λ being any real number. Prove that the points of division all lie on a curve similar to the two given curves except when they all happen to fall together.

CALCULUS.

When this issue was made up no solutions had been received for numbers 335, 337 to 340, 342, 348, and 350. Please give attention to these.

356. Proposed by B. F. FINKEL, Drury College.

A steel girder l feet long and w feet wide is moved on rollers along a passageway a feet wide and into a corridor at right angles to the passageway. How wide must the corridor be to just admit the girder?

357. Proposed by W. D. CAIRNS, Oberlin College.

A continuous variable represented by a point on a vertical line changes according to such a law that it is reduced to $1/m$ of its value on being moved a units upward, irrespective of the special position from which it is moved. Find the law of change, that is, the relation between the variable y and the height h of the variable point above a fixed point of the vertical line.

358. Proposed by C. N. SCHMALL, New York City.

About a given circle circumscribe the smallest parabola.

MECHANICS.

When this issue was made up no solutions had been received for numbers 271 to 275 inclusive, and 277 to 279 inclusive. Also 268 and 269, proposed in 1912, have not been solved. Please give attention to these.

288. Proposed by C. E. HORNE, Westminster College, Colorado.

Show that the tangential velocity of a projectile at any point of its path is equal to the velocity it would have acquired in falling, under the influence of gravitation alone, from the directrix to the point in question (Hulburt's *Calculus*, p. 220, ex. 10).

289. Proposed by C. N. SCHMALL, New York City.

A particle of elasticity e is projected with a velocity v at an angle of elevation ϕ from a point on a smooth horizontal plane. Show that after $2v \sin \phi / g(1 - e)$ seconds, it will cease to rebound and will move along the plane with a uniform velocity $v \cos \phi$.